arXiv:1609.08297v1 [q-bio.PE] 27 Sep 2016

Deterministic extinction by mixing in cyclically competing species

Cilie W. Feldager and Namiko Mitarai*
Center for Models of Life, Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, 2100 Copenhagen, Denmark.

Hiroki Ohta†
Niels Bohr International Academy / Center for Models of Life, Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, 2100 Copenhagen, Denmark.

(Dated: September 28, 2016)

We consider a cyclically competing species model on a ring with global mixing at finite rate, which corresponds to the well-known Lotka-Volterra equation in the infinite mixing rate. Within a perturbation analysis of the model from the infinite mixing rate, we provide analytical evidence that extinction occurs deterministically at sufficiently large but finite values of the mixing rate for any odd species numbers. Further, we discuss numerical results concerning the trajectories toward such deterministic extinction, including bifurcations, in more general cases.

PACS numbers: 87.23.Cc, 05.50.+q, 05.10.Gg

Introduction: One of the central interests in the field of theoretical ecology and game theory is to understand the mechanism of coexistence and extinction in interacting agents [1–3]. Among many kinds of models for interacting agents, N-species cyclic dominance has been studied intensely due to rich dynamics and possible long-term coexistence. Furthermore, its simplicity has facilitated theoretical studies [4–13], which could provide a fundamental viewpoint to understand more complicated cases observed in nature [14–17].

Dynamical behaviors of cyclic dominance between interacting agents may be described by the Lotka-Volterra (LV) equation or a cyclically competing species model on a lattice, where the former is supposed to be derived in the large size limit of the latter on a well-mixed condition. Specifically, the time evolution of the density P_α of a species α for $\alpha \in \{1, 2, \cdots, N\}$ in the LV equation is described as follows:

$$\dot{P}_\alpha = \frac{1}{2} P_\alpha (P_{\alpha_+} - P_{\alpha_-}), \tag{1}$$

where $\dot{P}_\alpha$ is the time derivative of P_α and $\alpha_+(\alpha_-)$ denoting a species forward (backward) next to α in the cyclic sense is a prey (predator) of species α (Fig. 1 left panel). This equation allows the coexistence of all the species where each density oscillates with an amplitude given by the initial condition; these trajectories are neutrally stable [1].

On the other hand, a cyclically competing species model on a lattice, where each species on a lattice stochastically invades its neighbors species along the rule of the cyclic dominance, shows different behaviors from those in the LV equation even qualitatively. For example, in the one-dimensional case, it has been proven that when $N \leq 4$, species at any sites are invaded after any time periods, showing slow coarsening, while when $N \geq 5$, each site reaches the final state, leading that the system gets frozen finally [7–10]. Although it has been poorly known how the behaviors of models depend on the dimensionality of the lattices, an approximation analysis rather suggests that the behaviors of the models do not approach those in the LV equation even if the spacial dimension is higher [11].



FIG. 1: (color online) A schematic description of the model. An invasion through the nearest neighbor (NN) occurs along the cyclic competetion. Individual species on randomly chosen two sites are exchanged at rate $v/2$.

As mentioned above, mixing in lattice models is supposed to bridge the gap between the two descriptions of cyclic ecosystems. However, if the system is not completely well-mixed, it could generally produce some macroscopic behaviors qualitatively different from those in the well-mixed condition. For example, one might consider mixing by local mobility, i.e., by locally exchanging the position of two species or also another mixing induced by a fluid flow, resulting in spatiotemporal density oscillations such as spirals [18, 19] or density oscillations with an increasing amplitude leading to extinction [20], respectively. Thus, such numerical works have already shown that mixing tends to induce oscillatory behaviors. However, how mixing makes a cyclic ecosystem lead to the LV equation is still not entirely clear. In this paper, we propose a simple model where one can extract analytical results, to some extent, on this topic.

In this letter, we study a cyclically competing species model on a ring by taking into account the effects of a global mixing at finite rate. By performing a perturbative analysis from the large mixing rate limit, we will

show that, at least, one species has to go *deterministically* extinct for cyclic competition of any odd species numbers even in the large size limit. This deterministic extinction is completely different from the extinction as a direct consequence of *stochasticity* [13]. It turns out that such deterministic extinction would be a key to understand why the relationship between the lattice models and the LV-equation for cyclic ecosystems was elusive.

Model: Let us denote by $\sigma_i \in \mathcal{S} \equiv \{1, \ldots, N\}$ with $N \geq 3$ a species at a site $i \in \{1, \ldots, L\}$ on a ring, i.e., a one-dimensional lattice with periodic boundary condition. Each species has one prey and one predator through a cyclic competition as shown in Fig. 1. Explicitly, we consider a continuous time Markov process, where the species at site i with $\sigma_i = \alpha$ invades the nearest neighbor sites $j \in \{i-1, i+1\}$ with rate $1/2$ only if $\sigma_j = \alpha_+$ where $\alpha_\pm \equiv \alpha \pm 1$ modulo N, making $\sigma_j = \alpha$ after invasion. In addition to this invading process, we introduce the exchange of two individual species on a uniformly chosen random pair of two sites i, j at rate $v/2$ as global mixing [21]: Namely, $(\sigma_i, \sigma_j) = (\alpha, \beta)$ with $\alpha, \beta \in \mathcal{S}$ is updated as $(\sigma_i, \sigma_j) = (\beta, \alpha)$ after mixing. Note that this mixing process does not change the population of each species in the system [22].

We define the n-point probability $P_{\{\alpha_k\}_{k=1}^n}(t)$ as the probability of a randomly chosen segment of n sites containing the sequence of species $\{\alpha_k\}_{k=1}^n$, in this order from the left to the right, with $\alpha_k \in \mathcal{S}$ for any integer k, $1 \leq k \leq n$ at time t. The time evolution of the one-point probability is exactly described by the two-point probability as

$$\dot{P}_\alpha = \frac{1}{2}(P_{\alpha,\alpha_+} + P_{\alpha_+,\alpha} - P_{\alpha_-,\alpha} - P_{\alpha,\alpha_-}). \tag{2}$$

In the limit of $L \to \infty$ and $v \to \infty$, one can easily find that this becomes the LV equation (1). Thus, in general, the evolution equation of $(n-1)$-point probability is exactly described by only n-point probability [11]. Note that Eq. (2) does not explicitly depend on the mixing rate v due to the conservation of the populations under mixing, while explicit contributions from mixing arise in $\dot{P}_{\{\alpha_k\}_{k=1}^n}$ for any $n \geq 2$.

Perturbative approach to deterministic extinction: We consider a perturbation approach from the infinite mixing rate where the limit of $L \to \infty$ is taken first. In order to obtain a closed set of evolution equations, it is a key to realize the following relation [23]:

$$P_{\{\alpha_k\}_{k=1}^n} = \frac{P_{\{\alpha_k\}_{k=1}^{n-1}} P_{\{\alpha_k\}_{k=2}^n}}{P_{\{\alpha_k\}_{k=2}^{n-1}}} + \mathcal{O}(v^{-(n-1)}), \tag{3}$$

where the error $\mathcal{O}(v^{-(n-1)})$, in the worst case, comes from the fact that species α_1 at the most left site never influences species α_n at the most right site unless α_1 succeed to invade all species to finally invade α_n through the segment of $(n-1)$-sites, where the probability that one invasion occurs before mixing to make predator α_1 away is $\mathcal{O}(1/v)$.

Using Eq. (3) with $n = 3$, we obtain the following equation for any $\alpha, \beta \in \mathcal{S}$:

$$\begin{aligned}\dot{P}_{\alpha,\beta} = \frac{1}{2}\Bigg(&- \delta_{\beta,\alpha_+} \frac{P_{\alpha,\beta}}{P_\beta} \sum_{\gamma \in \mathcal{S}\setminus\{\beta\}} P_{\beta,\gamma} - \frac{P_{\alpha,\beta} P_{\beta,\beta_-}}{P_\beta} \\ &- \frac{P_{\alpha_-,\alpha} P_{\alpha,\beta}}{P_\alpha} - \delta_{\alpha,\beta_+} \frac{P_{\alpha,\beta}}{P_\alpha} \sum_{\gamma \in \mathcal{S}\setminus\{\alpha\}} P_{\gamma,\alpha} \\ &+ \frac{P_{\alpha,\alpha_+} P_{\alpha_+,\beta}}{P_{\alpha_+}}(1 - \delta_{\beta,\alpha_+}) + \frac{P_{\alpha,\beta_+} P_{\beta_+,\beta}}{P_{\beta_+}}(1 - \delta_{\alpha,\beta_+}) \\ &+ \delta_{\alpha,\beta}(P_{\alpha,\alpha_+} + P_{\beta_+,\beta})\Bigg) \\ &+ 2v\,(P_\alpha P_\beta - P_{\alpha,\beta}) + \mathcal{O}(v^{-2}).\end{aligned} \tag{4}$$

Thus, Eq. (2) and Eq. (4) give a closed set of equations, allowing us to calculate exactly P_α and $P_{\alpha,\beta}$ within the errors of $\mathcal{O}(v^{-2})$.

We now move on to introduce an indicator:

$$\eta(t) \equiv \min_{\alpha \in \mathcal{S}} P_\alpha(t), \tag{5}$$

in order to discuss the extinction of, at least, one species, which is defined as $\eta \to 0$ as $t \to \infty$. Indeed, instead of directly focusing on η, it turns out that it is convenient to consider a quantity representing the intrinsic cyclic symmetry:

$$E(t) \equiv \prod_{\alpha \in \mathcal{S}} P_\alpha(t), \tag{6}$$

because $E = 0$ corresponds to $\eta = 0$ and thus one may conclude that extinction occurs by $E \to 0$ as $t \to \infty$. Note that one can easily find that E is a conserved quantity in the limit of $v \to \infty$ by analyzing directly the LV equation (1) [1].

Let us estimate $\dot{E}$ within $\mathcal{O}(v^{-2})$ by expanding $P_{\alpha,\beta}$ in the following way:

$$P_{\alpha,\beta} = P_\alpha P_\beta + \Delta_{\alpha,\beta} v^{-1} + \mathcal{O}(v^{-2}), \tag{7}$$

where the order of $\Delta_{\alpha,\beta}$ is $\mathcal{O}(v^0)$ due to relation (3) with $n = 2$. After obtaining the explicit expression of $\Delta_{\alpha,\beta}$ by substituting Eq. (7) into Eq. (4), one may finally obtain

$$\dot{E} \;=\; -\frac{1}{4} E \sum_{\alpha \in \mathcal{S}}^{N-1} \left(P_{\alpha_-} - P_{\alpha_+}\right)^2 v^{-1} + \mathcal{O}(v^{-2}). \tag{8}$$

On the assumption that this perturbation series has a finite converging radius in terms of v^{-1}, let us elaborate on what Eq. (8) means in terms of extinction. Indeed, $\dot{E}$ is always non-positive and may get zero only if the state is extinct with $E = 0$ or balanced with $P_{\alpha_-} = P_{\alpha_+}$ for any $\alpha \in \mathcal{S}$. Further, E takes the maximum value N^{-N} when the state is symmetric with $P_\alpha = 1/N$ for any $\alpha \in \mathcal{S}$. If the species number is odd, the symmetric condition corresponds to the balanced condition. Therefore, we



FIG. 2: (color online) (a) The maximum real part of the eigenvalues λ at the extinct fixed points for $N = 3$. The imaginary part is shown in the inset. (b) $P_\alpha(t)$ computed by Eqs. (2) and (4) for $v = 1$ from the symmetric initial condition with a perturbation.

may conclude that for any odd species numbers, $E \to 0$ as $t \to \infty$ leading to $\eta \to 0$ as $t \to \infty$ unless the initial state is precisely symmetric. If species number is even, $\dot{E} = 0$ might be achieved at also the asymmetric balanced state with $E \notin \{N^{-N}, 0\}$. This implies the possibility for any even species numbers that $E \to C$ with $0 < C < N^{-N}$. Nevertheless, we conjecture that such asymmetric balanced states would be not an attractor, because we could not find any such trajectories to reach those by both solving directly Eq. (4) and Eq. (7), and Monte Carlo (MC) simulations, which will be discussed below in more detail.

Extinction trajectories and bifurcation: We next discuss through which trajectory the system approaches the extinction $\eta \to 0$, in particular, by employing the linear stability analysis of Eq. (2) and (4) at both extinct, meaning $P_\alpha P_{\alpha_+} = 0$ for any $\alpha \in \mathcal{S}$, and symmetric, meaning $P_\alpha = 1/N$ for any $\alpha \in \mathcal{S}$, fixed points with $\dot{P}_\alpha = 0, \dot{P}_{\alpha,\beta} = 0$ for any $\alpha, \beta \in \mathcal{S}$ [24]. Note that such a linear stability analysis for finite v should be referred to as an approximation, but it would give us insightful information as found below. Hereafter, we call the eigenvalue with the largest real part among all the eigenvalues as the dominant eigenvalue and the corresponding eigenvector as the dominant eigenvector. If the real part of the dominant eigenvalue is positive, the dominant eigenvector is called also the most unstable eigenvector. Let us discuss the cases of $N = 3$, $N = 4$, and also $5 \le N \le 8$ in this order below.

In the case of 3-species competition ($N = 3$), there are three extinct fixed points with $P_\alpha = 1$, $P_{\alpha_+} = P_{\alpha_-} = 0$ for any $\alpha \in \mathcal{S}$. The linear stability analysis indicates that the dominant eigenvalue is positive at any finite v as shown in Fig. 2a. Thus, the extinct fixed points turn out to be saddles. Therefore, one consistent way to realize the trajectory to extinction satisfying $\eta \to 0$ is a heteroclinic cycle connecting such unstable extinct fixed points. As shown in Fig. 2b, by solving numerically Eq. (2) and Eq. (4) from the initial conditions close to the symmetric fixed point, we indeed observed such heteroclinic cycles.



FIG. 3: (color online) (a) The maximum real part of eigenvalues at the extinct fixed point for $N = 4$ as a function of p. (b) Eigenvalues λ at the symmetric fixed point with $N = 4$. The top panel shows the two eigenvalues with the largest and the second largest real part, and the bottom panel shows their respective imaginary parts. (c) $P_\alpha(t)$ computed by Eqs. (2) and (4) with $N = 4$ from the symmetric initial condition with a perturbation for $v = 0.02$. (d) $P_\alpha(t)$ computed by Eqs. (2) and (4) with $N = 4$ from the symmetric initial condition with a perturbation for $v = 1$. (e-f) MC simulations: The species densities with $L = 9000$ (e,f) and the spatiotemporal plot with $L = 1000$ (g,h) for $v = 0.02$ (e,g) and $v = 1$ (f,h), where x is site number.

This kind of trajectories was also observed by looking at the species densities in MC simulations at rather small v. This implies that mixing always causes the deterministic extinction for $N = 3$ even with small v, leading to the heteloclinic cycle.

In the case of 4-species competition, there are two connected regions parametrized by a real number $p \in [0, 1]$, consisting of extinct fixed points $P_\alpha = p$, $P_{\alpha_{++}} = 1 - p$, and $P_{\alpha_+} = P_{\alpha_-} = 0$ for any $\alpha \in \mathcal{S}$, that allows survival

of two species. As shown in Fig. 3a, we found that there is a region in the phase space, close to $p = 1/2$, where only the dominant eigenvalue is zero, meaning that all the rest eigenvalues have negative real parts. Since the direction along the connected extinct fixed points in the phase space is obviously neutrally stable, corresponding the zero eigenvalue, these extinct fixed points around $p = 1/2$ compose an attractor. As v is increased, the extinct fixed points for smaller p get to lose its local stability, leading that this extinct attractor disappears in the limit of $v \to \infty$. Oppositely, as v is decreased from a sufficiently large value, we found a bifurcation at $v = v_c \approx 0.19$, where the most unstable eigenvector at the symmetric fixed point discontinuously changes from an oscillatory mode to a non-oscillatory mode, as shown in Fig. 3b. Consistent with these results, the following were found by solving numerically Eq. (2) and Eq. (4) from the initial conditions close to the symmetric fixed point. For $v = 1 > v_c$, the trajectories seem to end up with heteroclinic cycles as shown Fig. 3d, implying that the basin of the extinct attractor around $p = 1/2$ is far from the symmetric fixed point in this case. However, for $v = 0.02 < v_c$, there are trajectories that fall into the extinct attractor as shown in Fig. 3c. Thus, at the bifurcation point, from the viewpoint of the phase space, the symmetric fixed point seems to collapse to a boundary of the basin of the extinct attractor. We also observed these two different types of trajectories by MC simulation as shown in Fig. 3(e-f).

We performed the linear stability analysis at the symmetric fixed point for also larger N. In the cases of $N = 5, 6, 7, 8$, we found discontinuous changes in the most unstable eigenvector at certain values of v. In the case of even N, as observed in 4 species, there is an eigenvector with zero imaginary part for any finite v, that becomes the most unstable eigenvector through a bifurcation as v is decreased. In the case of odd N, the dominant eigenvector has non-zero imaginary part for any finite v and the oscillation amplitude dramatically changes at the bifurcation point. Note that for $N = 3$, we did not find any discontinuous changes in the most unstable eigenvector. Finally, for general N as far as we have studied, the real part of the dominant eigenvalue is non-negative and approaches zero in the limit of both $v \to 0$ and $v \to \infty$. Thus, from the viewpoint of this instability of the symmetric fixed point, the limits of $v \to 0$ and $v \to \infty$ are rather singular.

Concluding remarks: In this letter, we have studied a cyclically competing species model on a ring with a global mixing. With a perturbative analysis combined with numerical methods, we have clarified that the deterministic extinction occurs for finite v and its singular appearance from the limits of $v \to 0$ and $v \to \infty$ is a key to bridge the gap between two descriptions of the LV equation and lattice models.

The global mixing studied in this letter is not only of theoretical interest. For example, a bacterial system consisting of toxin-producing, sensitive, and immune bacteria strains can form a cyclic competition and the mixing process in a liquid culture could be effectively global because of a vigorous mixing [14, 15]. The present results in this paper imply that the effects of deterministic extinction could be rather widely observed in experiments because of the difficulties to realize precisely a well-mixed condition in reality. In experiments, we expect that the effects from deterministic extinction are observed when each species density is sufficiently far from zero, and eventually one species goes extinct because of intrinsic stochasticity in the system when the minimum species density gets close to zero compared to the strength of stochasticity.

Lastly, let us mention the universality of the deterministic extinction in terms of different descriptions. Indeed, deterministic extinction for 3-species cyclic competition has been observed widely in the description by an ordinary differential equation with logistic growth and death [5], in a lattice model for competing species under a flow at fast mixing [20], and also in the framework of the continuous limit described by a partial differential equation of populations under a convecting turbulent flow at fast mixing [25]. Such observations of deterministic extinction in different descriptions imply the universality of deterministic extinction in cyclic systems with fast mixing, to some extent. Thus, we hope that the obtained results in this paper could provide a fundamental viewpoint to develop our understanding of mixing-induced deterministic extinction in even more general ecosystems.

The authors thank E. Frey for useful discussions on the topic related to Ref. [25]. This work is supported by the Danish National Research Foundation.

* Electronic address: mitarai@nbi.dk
† Electronic address: hirokiohta@jamstec.go.jp

[1] J. Hofbauer and K. Sigmund, *The theory of evolution and dynamical systems: mathematical aspects of selection* (Cambridge University Press, 1988).
[2] G. Szabó and G. Fath, Phys. Rep. **446**, 97 (2007).
[3] A. Szolnoki, M. Mobilia, L.-L. Jiang, B. Szczesny, A. M. Rucklidge, and M. Perc, J. R. Soc. Interface **11**, 20140735 (2014).
[4] Y. Itoh, Proc. Japan Acad. **47**, 854 (1971).
[5] R. M. May and W. J. Leonard, SIAM J. Appl. Math. **29**, 243, (1975).
[6] K. Tainaka, J. Phys. Soc. Jpn. **57**, 2588 (1988).
[7] M. Bramson and D. Griffeath, Ann. Prob. **17**, 26 (1989).
[8] R. Fisch, Ann. Prob. **20**, 1528 (1992).
[9] L. Frachebourg, P. L. Krapivsky, and E. Ben-Naim, Phys. Rev. E **54**, 6186 (1996).
[10] L. Frachebourg, P. L. Krapivsky, and E. Ben-Naim, Phys. Rev. Lett. **77**, 2125 (1996).
[11] L. Frachebourg and P. L. Krapivsky, J. Phys. A: Math. Gen. **31**, L287 (1998).

[12] R. Durrett and S. Levin, Theor. Popul. Biol. **53**, 30 (1998).
[13] T. Reichenbach, M. Mobilia, E. Frey, Phys. Rev. E **74**, 051907 (2006).
[14] T. L. Czárán, R. F. Hoekstra, and L. Pagie, Proc. Natl. Acad. Sci. **99**, 786 (2002).
[15] B. Kerr, M. A. Riley, M. W. Feldman, and B. J. M. Bohannan, Nature **418**, 171 (2002).
[16] J. Mathiesen, N. Mitarai, K. Sneppen, and A. Trusina, Phys. Rev. Lett. **107**, 188101 (2011).
[17] Z. Frentz, S. Kuehn, and S. Leibler, Phys. Rev. X **5**, 041014 (2015).
[18] T. Reichenbach, M. Mobilia, E. Frey, Phys. Rev. Lett. **99**, 238105 (2007).
[19] T. Reichenbach, M. Mobilia, E. Frey, Nature **448**, 1046 (2007).
[20] G. Károlyi, Z. Neufeld, and I. Scheuring, J. Theor. Biol. **236**, 12 (2005).
[21] A. R. Akhmetzhanov, L. Worden, and J. Dushoff, Phys. Rev. E **88**, 012816 (2013).
[22] As a mixing process, one can consider a copy-paste process where a species at a randomly chosen site invades a species at another randomly chosen site with rate v. In this case, the main results in this paper are unchanged.
[23] Note that the case with $n = 3$ was used in Ref. [11] for $v = 0$.
[24] We computed $P_{\alpha,\beta}$ at the symmetric fixed points by solving numerically Eqs. (2) and (4) from the initial condition where $P_\alpha = 1/N, P_{\alpha,\beta} = \delta_{\alpha,\beta}/N$, where $\delta_{a,b}$ is the kronecker delta function. By using this initial condition with no invasions, one can avoid coarsening effects causing slow relaxations. At the extinct fixed points, one can obtain the exact relation $P_{\alpha,\beta} = P_\alpha P_\beta$.
[25] D. Grošelj, F. Jenko, and E. Frey, Phys. Rev. E **91**, 033009 (2015).